

## The surgical treatment of vertigo

N. Deggouj, S. Castelein and M. Gersdorff

Academic Hospital Saint-Luc, ENT Department, Université catholique de Louvain, Brussels, Belgium

**Key-words.** Surgery; vertigo; labyrinthectomy; neurectomy; fistula

**Abstract.** *The surgical treatment of vertigo.* We present surgical treatments for vertigo targeting symptoms and aetiology.

The choice of treatment should be guided by the symptoms, the supposed aetiological factors and the age of the subject. Surgery is rarely indicated in the majority of the patients with vertigo.

Surgical solutions should be proposed for cases of chronic incapacitating vertigo that are refractory to medical or physical treatment.

The surgical procedures possible today target aetiological factors or attempt to cancel information coming from the vestibular organ. We will refer to the latter as "symptomatic treatment".

### A. Symptomatic surgical treatment

These treatments destroy all or part of the labyrinth, or disconnect it from the vestibular centres.

#### 1. Labyrinthectomy

##### a) Total labyrinthectomy

This procedure may be either chemical or surgical. The intratympanic application of Gentamycin is the gold standard for the treatment of intractable chronic vertigo of the kind found in Ménière's disease.<sup>1-4</sup> Ideally, the aminoglycosides are administered in the round window niche through a paracentesis, a transtympanic tube or sometimes through a pump (not available now). The easiest solution today is to administer it using the Spongostan of a Silverstein tube (Micromedics Inc. Minnesota 5510 USA). This treatment results in good control of the vertigo (>80%) but it may adversely affect hearing. Surgical labyrinthectomy should be avoided. Otolologists

should aim to preserve cochlear morphology so that cochlear implantation remains possible at a later stage. The destruction of the diseased end organ will eliminate vertigo symptoms but also destroys hearing. This procedure should be reserved for ears with significant hearing loss in old patients.

##### b) Localised labyrinthectomy

The plugging of a semi-circular canal is an effective treatment with >90% success in vertigo control in cases of intractable benign paroxysmal positional vertigo (BPPV).<sup>5-7</sup> It usually involves the posterior canal and sometimes the horizontal. Incomplete occlusion of the canal may result in recurrence of BPPV and in a catiogenic fistula.<sup>8</sup>

#### 2. Vestibular neurectomies and neurectomies

##### a) Singular neurectomy

This approach is proposed for chronic disabling refractory BPPV. The neurectomy is carried

out in the ampullary recess of the posterior semi-circular canal. This technique provides complete relief from BPPV in 96% of subjects.<sup>9</sup> Sensorineural hearing loss is a complication in 3-4%. A positive fistula response may be present for a few months in some patients.

##### b) Vestibular neurectomy

A vestibular neurectomy in the pontocerebellar angle allows for the disconnection of the vestibular end organ from the vestibular nuclei and the cerebellum. It usually results in a central vestibular compensatory reorganisation, allowing for the recovery of functional equilibrium. This compensation can happen despite the persistent evolutive disease in the ear or the nerve. It is less efficient in the elderly and so this will not be the first-choice surgical option for them.

A vestibular neurectomy results in improvements or the resolution of vertigo in >90%.<sup>10-12</sup> The retrosigmoid approach is simpler, safer and more reliable than the middle or retrolabyrinthine

approaches.<sup>11</sup> The retrosigmoid approach has been used to associate neurectomy and vascular decompression.<sup>13</sup> A recurrence of vertigo after surgery can be observed. This may be due to an incomplete neurectomy or, more frequently, to the passage of vestibular neural fibres into the preserved cochlear nerve.<sup>14</sup>

## B. Aetiological surgical treatments

The aim of these treatments is to correct the supposed aetiological factors of vertigo.

### 1. Treatments acting on the pressure in the ear

Those treatments were developed historically to cure or to improve an endolymphatic hydrops (EH), which was presumed to explain the symptoms. The first decompressive surgical treatments consisted of sacculotomy or cochleo-sacculotomy. They were intended to reduce EH by drainage and therefore improve the symptoms.

They were found to be effective for vertigo in 80%.<sup>15,16</sup> However, the risk of associated deterioration in hearing was higher: 20%.<sup>16</sup>

Decompressive surgery procedures involving the endolymphatic sac continue to be used by many teams.<sup>17-26</sup>

In fact, all these decompressing sac surgeries have been found to be unspecific.<sup>16,19,21</sup> They have the same positive effect on vertigo improvement as a simple mastoidectomy<sup>19</sup> on transtympanic ventilation tube in humans.<sup>15,16</sup> In animal models of EH, the cochleo-sacculotomy was not found to have any effect on the induced EH.<sup>15</sup>

A severe permanent hearing loss may occur in 20% of patients

after sacculotomy and cochleo-sacculotomy.<sup>16</sup>

A transtympanic tube seems preferable to sac surgery in Ménière's disease.

This tube makes it possible to apply low-intensity alternating pressure. The Meniett device seems to be a promising and efficacious treatment for Ménière's subjects with vertigo uncontrolled by other treatments.<sup>27</sup>

### 2. Treatment of fistula

Labyrinthine fistulae may be primitive or secondary to an inflammatory process in the middle ear (cholesteatoma, granulomatosis); to surgery (cholesteatoma, cochlear implantation otosclerosis); to traumatism; to malformations (Mondini). In the case of symptomatic fistulae, there should be a surgical exploration of the middle ear and the fistula should be packed with periosteum muscle or other tissues.<sup>28-30</sup>

The superior semicircular canal dehiscence is a special type of fistula.<sup>31-33</sup> It must be treated using a combined otological and neurosurgical procedure through the middle cranial fossa.

### 3. Other

Some cases of Tullio's phenomenon are due to abnormal mobility of the stapes or other ossicles. The hypermobility may be stabilised by placing cartilage chips beside the crurae of the stapes.<sup>34</sup> Abnormal scar tissue around the ossicles may be resected during a surgical exploration of the middle ear.

## Conclusion

Rapid surgical treatment is indicated when a aetiological factor

may be safely corrected. In other cases, the surgical option should be reserved for incapacitating vertigo that does not respond to medical or physical treatment. The choice of treatment must be guided by the symptoms, experience and the age of the subjects.

## References

1. Cohen Kerem R, Kisilevsky V, Einarson TR, Kozar E, Koren G, Rutka JA. Intratympanic gentamicin for Ménière's disease: a meta-analysis. *Laryngoscope*. 2004;114: 2085-2091.
2. Hoffer ME, Kopke RD, Weisskopf P, et al. Use of the round window microcatheter in the treatment of Ménière's disease. *Laryngoscope*. 2001;111: 2046-2049.
3. Seidman M. Continuous gentamicin therapy using an IntraEAR microcatheter for Ménière's disease: a retrospective study. *Otolaryngol Head Neck Surg*. 2002;126:244-256.
4. Van de Heyning PH, Wuyts F, Boudewyns A. Surgical treatment of Ménière's disease. *Curr Opin Neurol*. 2005;18:23-28.
5. Crossland G, De R, Axon P. Ear advanced otosclerosis and intractable benign paroxysmal positional vertigo treated with combined cochlear implantation and posterior semicircular canal occlusion. *J Laryngol Otol*. 2004;118:302-304.
6. Horii A, Imai T, Mishiro Y, et al. Horizontal canal type BPPV: bilaterally affected case treated with canal plugging and Lempert's maneuver. *ORL J Otorhinolaryngol Relat Spec*. 2003;65:366-369.
7. Agrawal SK, Parnes LS. Human experience with canal plugging. *Ann N Y Acad Sci*. 2001;942:300-305.
8. Rizvi SS, Gauthier MG. Unexpected complication of posterior canal occlusion surgery for benign paroxysmal positional vertigo. *Otol Neurotol*. 2002;23:938-940.
9. Gacek RR, Gacek MR. Results of singular neurectomy in the posterior ampullary recess. *ORL J Otorhinolaryngol Relat Spec*. 2002;64:397-402.

10. Pappas DG Jr, Pappas DG Sr. Vestibular nerve section: long-term follow-up. *Laryngoscope*. 1997;107:1203-1209.
11. Magnan J, Bremond G, Chays A, Gignac D, Florence A. Vestibular neurectomy by retrosigmoid approach: technique, indications and results. *Am J Otol*. 1991;12:101-104.
12. Pareschi R, Destito D, Falco Raucci A, Righini S, Colombo S. Posterior fossa vestibular neurectomy as primary surgical treatment of Meniere's disease: a re-evaluation. *J Laryngol Otol*. 2002;116:593-596.
13. Zini C, Mazzoni A, Gandolfi A, Sanna M, Pasanisi E. Retrolabyrinthine versus middle fossa vestibular neurectomy. *Am J Otol*. 1988;9:448-450.
14. Lehnen N, Aw ST, Todd MJ, Halmagyi GM. Head impulse test reveals residual semicircular canal function after vestibular neurectomy. *Neurology*. 2004;62:2294-2296.
15. Montandon P, Guillemin P, Häusler R. Prevention of vertigo in Ménière's syndrome by means of transtympanic ventilation tubes. *ORL J Otorhinolaryngol Relat Spec*. 1988;50:377-381.
16. Häusler R, Guillemin P, Montandon P. Surgical treatment of Ménière's disease by sacculotomy, cochleo-sacculotomy and transtympanic aerators [in French]. *Rev Laryngol Otol Rhinol*. 1991;112:149-152.
17. Arenberg IK, Stahle J, Wilbrand H, Newkirk JB. Unidirectional inner ear valve implant for endolymphatic sac surgery in Meniere's disease. *Arch Otolaryngol*. 1978;104:694-704.
18. Kato BM, LaRouere MJ, Bojrab DI, Michaelides EM. Evaluating quality of life after endolymphatic sac surgery: The Ménière's Disease Outcomes Questionnaire. *Otol Neurotol*. 2004;25:339-344.
19. Thomsen J, Bretlau P, Tos M, Johnsen NJ. Placebo effect in surgery for Meniere's disease. A double-blind, placebo-controlled study on endolymphatic sac shunt surgery. *Arch Otolaryngol*. 1981;107:271-277.
20. Thomsen J, Bretlau P, Tos M, Johnsen NJ. Endolymphatic sac-mastoid shunt surgery. A nonspecific treatment modality? *Ann Otol Rhinol Laryngol*. 1986;95:32-35.
21. Thomsen J, Kerr A, Bretlau P, Olsson J, Tos M. Endolymphatic sac surgery: why we do not do it. The non-specific effect of sac surgery. *Clin Otolaryngol Allied Sci*. 1996;21:208-211.
22. Huang TS. Endolymphatic sac surgery for Meniere's disease: experience with over 3000 cases. *Otolaryngol Clin North Am*. 2002;35:591-606.
23. Welling DB, Pasha R, Roth LJ, Barin K. The effect of endolymphatic sac excision in Ménière's disease. *Am J Otol*. 1996;17:278-282.
24. Kitahara T, Takeda N, Mishiro Y, et al. Effects of exposing the opened endolymphatic sac to large doses of steroids to treat intractable Meniere's disease. *Ann Otol Rhinol Laryngol*. 2001;110:109-112.
25. Moody-Antonio S, House JW. Hearing outcome after concurrent endolymphatic shunt and vestibular nerve section. *Otol Neurotol*. 2003;24:453-459.
26. Bertran JM, Alvarez de Cozar F. Results of surgery of the endolymphatic sac [in Spanish]. *Acta Otorhinolaryngol Esp*. 1999;50:179-183.
27. Gates GA, Green JD Jr, Tucci DL, Telian SA. The effects of transtympanic micropressure treatment in people with unilateral Meniere's disease. *Arch Otolaryngol Head Neck Surg*. 2004;130:718-725.
28. Kusuma S, Liou S, Haynes DS. Disequilibrium after cochlear implantation caused by a perilymph fistula. *Laryngoscope*. 2005;115:25-26.
29. Fox EJ, Balkany TJ, Arenberg IK. The Tullio phenomenon and perilymph fistula. *Otolaryngol Head Neck Surg*. 1988;98:88-89.
30. Manolidis S. Complications associated with labyrinthine fistula in surgery for chronic otitis media. *Otolaryngol Head Neck Surg*. 2000;123:733-737.
31. Ramsey MJ, McKenna MJ, Barker FG 2nd. Superior semicircular canal dehiscence syndrome. Case report. *J Neurosurg*. 2004;100:123-124.
32. Minor LB. Superior canal dehiscence syndrome. *Am J Otol*. 2000;21:9-19.
33. Deggouj N, Castelein S, Gersdorff M. Tullio's phenomenon. *B-ENT*. 2008; suppl. 8:27-28.
34. Jahnke K, Dost P. Surgical treatment of acoustically-induced vertigo (Tullio phenomenon) [in German]. *Laryngorhinootologie*. 1999;78:394-396.

Naïma Deggouj, M.D.  
Academic Hospital Saint-Luc  
ENT Department  
Université catholique de Louvain  
avenue Hippocrate 10  
B-1200 Brussels, Belgium  
Tel.: 32.2.764.19.49.  
Fax: 32.2.764.89.35.  
E-mail: Naima.Deggouj@uclouvain.be